

sphalm. 1965.

Bibliography: Körn. in Mart., Fl. Bras. 3 (1): 456—458. 1863; Ruhl. in Engl., Pflanzenreich 13 (4-30): 254 & 293. 1903; Alv. Silv., Fl. Mont. 1: 419. 1928; Moldenke, Known Geogr. Distrib. Erioc. 19, 51, & 59. 1946; Moldenke, Phytologia 2: 498. 1948; Moldenke, Known Geogr. Distrib. Verbenac., [ed. 2], 92 & 213. 1949; Moldenke, Phytologia 4: 326. 1953; Angely, Fl. Paran. 10: 15. 1957; Moldenke, Résumé 108, 117, 326, 352, & 492. 1959; Angely, Fl. Paran. 16: 77 (1960) and 17: 24. 1961; Angely, Fl. Anal. Paran., ed. 1, 202. 1965; Moldenke, Résumé Suppl. 18: 14. 1969; Moldenke, Fifth Summ. 1: 175 & 187 (1971) and 2: 587, 637, 964, & 968. 1971; Angely, Fl. Anal. & Fitogeogr. Est. S. Paulo, ed. 1, 1163 & Ind. 28. 1972. [to be continued]

NOTES ON NEW AND NOTEWORTHY PLANTS. CIV

Harold N. Moldenke

SYNGONANTHUS TENUIS var. *MINOR* Moldenke, var. nov.

Haec varietas a forma typica speciei statura humilior pedunculis 4—8 cm. altis capitulis minoribus recedit.

This variety differs from the typical form of the species in its lower stature, the peduncles only 4—8 cm. long, and the smaller heads.

The type of the form was collected by Julian A. Steyermark, G. C. K. Dunsterville, and E. Dunsterville (no. 113217) at "cumbre de Cerro Guaiquinima, Salto del Río Szczerbanari (Río Carapo), 1--2 km. río arriba del Salto Szczerbanari, lat. 5°44'4" N., long. 63°31'8" W., parte central del cerro", at 750 m. altitude, Bolívar, Venezuela, between January 20 and 25, 1977, and is deposited in my personal herbarium.

VERBENA RIGIDA f. *ALBA* (Trivetts) Moldenke, stat. nov.

Verbena venosa alba Trivetts, Cat. Suppl. 1933 Novelty List 6. 1933.

VERBENA RIGIDA f. *LILACINA* (Benary & Bodger) Moldenke, stat. nov.

Verbena venosa lilacina Benard & Bodger ex Harrow, Journ. Roy. Hort. Soc. 61: 401. 1935.

VITEX PEDUNCULARIS f. juv. *ROXBURGHIANA* (C. B. Clarke) Moldenke, stat. nov.

Vitex peduncularis var. *roxburghiana* C. B. Clarke in Hook. f., Fl. Brit. India 4: 587. 1885.